


Current Biology

Multi-sensor tracking reveals the extent of nocturnal migration in landbirds

Graphical abstract



Authors

Paul Dufour, Raphaël Nussbaumer, Paul Doniol-Valcroze, ..., Tianhao Zhao, Barbara Helm, Martins Briedis

Correspondence

paul.dufour80@gmail.com

In brief

Dufour et al. combine multi-sensor geolocator data from 411 individuals across 56 landbird species to quantify nocturnal and diurnal migratory flight. They show that nocturnal migration is the dominant strategy among small landbirds, is phylogenetically conserved, and that several species previously considered diurnal migrants in fact fly exclusively at night.

Highlights

- Multi-sensor geolocators are uniquely suited to quantify flight timing in landbirds
- Nocturnal flights dominate (82%) across 56 small- to medium-sized landbird species
- Departures and landing times cluster around civil dusk and dawn
- Nocturnal flight timing shows moderate to strong phylogenetic signal



Report

Multi-sensor tracking reveals the extent of nocturnal migration in landbirds

Paul Dufour,^{1,48,*} Raphaël Nussbaumer,¹ Paul Doniol-Valcroze,² Yann Rime,³ Peter Adamík,⁴ Susanne Åkesson,⁵ Raphaël Arlettaz,⁶ Arnaud G. Barras,⁷ Nyambayar Batbayar,⁸ Ian Beggs,⁹ Fraser Bell,^{10,11} Alberto Benito Ruiz,¹² Staffan Bensch,⁵ Pierrick Bocher,¹³ Than Boves,¹⁴ Jesper Brodersen,⁵ Malcolm Burgess,^{10,11,15} Lars Burnus,¹⁶

(Author list continued on next page)

¹Bird Migration Unit, Swiss Ornithological Institute, Sempach 6204, Switzerland

²CESCO, UMR7204 MNHN-CNRS-Sorbonne Université, Paris 75005, France

³FitzPatrick Institute of African Ornithology, University of Cape Town, Rondebosch 7701, South Africa

⁴Department of Zoology, Faculty of Science, Palacký University, Olomouc 772 00, Czech Republic

⁵Department of Biology, Lund University, Lund 223 62, Sweden

⁶Division of Conservation Biology, Institute of Ecology and Evolution, University of Bern, Bern 3000, Switzerland

⁷Regional Office Valais, Swiss Ornithological Institute, Sion 1950, Switzerland

⁸Wildlife Science and Conservation Center of Mongolia, Ulaanbaatar 14200, Mongolia

⁹Skokholm Island Bird Observatory, Pembrokeshire, UK

¹⁰Center for Ecology and Conservation, University of Exeter, Penryn TR10 9FE, UK

¹¹PiedFly.Net, Yarnar Wood, Newton Abbot, Devon TQ13 9LJ, UK

¹²Sociedad Española de Ornitología, SEO/BirdLife, Madrid 28001, Spain

¹³University of La Rochelle, La Rochelle 17000, France

¹⁴Department of Biological Sciences, Arkansas State University, Jonesboro, AR 72401, USA

¹⁵RSPB Centre for Conservation Science, Sandy SG19 2DL, UK

¹⁶Institute of Avian Research "Vogelwarte Helgoland", Wilhelmshaven 26386, Germany

(Affiliations continued on next page)

SUMMARY

Most bird species are diurnal but drastically change their diel cycle of activity to migrate at night. Nocturnal migration has been documented using different methods (e.g., experiments, radar and radio tracking, and acoustic monitoring), but accurately quantifying the proportion of nocturnal versus diurnal flight at the individual and species levels, and understanding how this behavior evolved across the avian tree, has remained methodologically challenging.^{1–3} Such uncertainty is not only of theoretical importance but also limits our ability to mitigate conservation threats, particularly from light pollution and building collisions.^{4–6} With multi-sensor geolocators recording light, barometric pressure, and activity,^{7,8} we reconstructed high-resolution migratory trajectories^{9,10} for 411 individuals from 56 small- or medium-sized landbird species across four continents and measured the proportion of each flight that occurred during night or day. Our species-level quantification confirmed nocturnal migration as the dominant strategy among small landbirds, while also providing precise flight proportion estimates across a broad taxonomic sample and refining the classification of several species previously described as partial or facultative diurnal migrants. We found that birds initiated and ended migratory flights near civil dusk and dawn, thereby maximizing nocturnal travel. The phylogenetic signal we detected indicates that nocturnal migration is largely conserved within lineages. While nocturnal migration confers multiple advantages, the relative importance of the proposed drivers remains to be determined.^{11,12} Our study provides species-specific quantification of nocturnal migration, highlights tracking gaps across taxa and regions, and opens new avenues for studying the evolutionary, ecological, and sensory drivers of nocturnal migratory flights.

RESULTS AND DISCUSSION

Multi-sensor geolocators allow quantification of nocturnal flight timing

The nocturnal migration of otherwise diurnal birds has long fascinated scientists and remains an important, unresolved

question.¹³ It represents a unique exception to the general pattern of diel activity rhythm in diurnal animals, which are typically active during the day and inactive at night.^{14,15} The first studies of nocturnal migration relied on moon-watching, experimental observations, including the quantification of migratory restlessness (*Zugunruhe*), and ringing activities and were later



Violeta Caballero-Lopez,⁵ Pablo Capilla-Lasheras,^{1,17,18} Jocelyn Champagnon,¹⁹ Andrea Contina,²⁰ Greg Conway,²¹ Nathan W. Cooper,²² Batmunkh Davaasuren,⁸ Emily R. Donahue,¹⁴ Bryant C. Dossman,²³ Olivier Duriez,²⁴ Ruben Evens,²⁵ Frode Fossøy,²⁶ Katie J. Galletta,²⁷ Astrid Günther,¹⁶ Anders Hedenström,⁵ Wieland Heim,²⁸ Chris Hewson,²¹ Marc Illa,²⁹ Colin Jackson,³⁰ Frédéric Jiguet,² Erik I. Johnson,^{31,32} Matthew Johnson,³³ Diane Klement,³⁴ Sonam Kulkarni,¹⁶ Michiel Lathouwers,³⁵ Felix Liechti,³⁶ Miriam Liedvogel,¹⁶ Dave Lutgen,³⁷ Pablo Macías-Torres,⁵ Christoph M. Meier,¹ Jugdernamjil Nergui,⁸ Samuel Temidayo Osinubi,³ Gonzalo Pardo de Santayana,¹² Petr Procházka,³⁸ Jirí Reif,^{4,39} Garrett Rhyné,⁴⁰ Clark Rushing,³⁴ Sissel Sjöberg,⁵ Bård G. Stokke,²⁶ Stephanie Szarmach,^{22,41} Kasper Thorup,⁴² David Toews,⁴¹ Christopher M. Tonra,⁴³ Anders P. Tøttrup,⁴⁴ Jennifer Tyrrell,³⁴ Joanna B. Wong,¹ Alexander Worm,¹⁴ Joe Wynn,^{16,45} Elizabeth Yohannes,¹ Tianhao Zhao,⁴⁶ Barbara Helm,¹ and Martins Briedis^{1,47}

¹⁷Doñana Biological Station (CSIC), Seville 41013, Spain

¹⁸School of Biodiversity, One Health and Veterinary Medicine, University of Glasgow, Glasgow G12 8QQ, UK

¹⁹Tour du Valat, Research institute for conservation of Mediterranean wetlands, Arles 13200, France

²⁰School of Integrative Biological and Chemical Sciences, The University of Texas Rio Grande Valley, Brownsville, TX 78520, USA

²¹British Trust for Ornithology, The Nunnery, Thetford, Norfolk IP24, UK

²²Migratory Bird Center, Smithsonian's National Zoo and Conservation Biology Institute, Washington, DC 20008, USA

²³Department of Biology, Georgetown University, Washington, DC 20057, USA

²⁴CEFE, University Montpellier, CNRS, EPHE, IRD, Montpellier 34000, France

²⁵UCLouvain, Earth & Life Institute, Nox'alis Group, Louvain-la-Neuve 1348, Belgium

²⁶Norwegian Institute for Nature Research, Trondheim 7485, Norway

²⁷Natural Resources & Environmental Sciences Department, University of Illinois Urbana-Champaign, Urbana, IL 61801, USA

²⁸Institute of Biology and Environmental Sciences, University of Oldenburg, Oldenburg 26129, Germany

²⁹Institut Català d'Ornitologia, Barcelona, Catalunya 08024, Spain

³⁰A Rocha Kenya, Nairobi 00502, Kenya

³¹Louisiana State University AgCenter, Baton Rouge, LA 70803, USA

³²Audubon Delta, National Audubon Society, New Orleans, LA 70130, USA

³³Audubon South Carolina, National Audubon Society, Charleston, SC 29403, USA

³⁴Warnell School of Forestry and Natural Resources, University of Georgia, Athens, GA 30602, USA

³⁵Research Institute for Nature and Forest (INBO), Brussels 1000, Belgium

³⁶University of Basel, Basel 4051, Switzerland

³⁷Monitoring Unit, Swiss Ornithological Institute, Sempach 6204, Switzerland

³⁸Czech Academy of Sciences, Institute of Vertebrate Biology, Brno 603 65, Czech Republic

³⁹Institute of Environmental Studies, Faculty of Science, Charles University, Prague 128 00, Czech Republic

⁴⁰American Bird Conservancy, Charlottesville, VA 22903, USA

⁴¹Department of Biology, The Pennsylvania State University, University Park, PA 16802, USA

⁴²Center for Macroecology, Evolution and Climate, University of Copenhagen, Copenhagen 2100, Denmark

⁴³School of Environment and Natural Resources, The Ohio State University, Columbus, OH 43210, USA

⁴⁴Natural History Museum Denmark, University of Copenhagen, Copenhagen 1172, Denmark

⁴⁵School of Environmental Sciences, University of Liverpool, Liverpool L69 3BX, UK

⁴⁶Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, Groningen, 9747AG, the Netherlands

⁴⁷Lab of Ornithology, Institute of Biology, Faculty of Medicine and Life Sciences, University of Latvia, Riga, LV-1004, Latvia

⁴⁸Lead contact

*Correspondence: paul.dufour80@gmail.com

<https://doi.org/10.1016/j.cub.2026.06.030>

complemented by radar observations, acoustic monitoring, and thermal imaging.^{16–19} Radar studies have long confirmed that nocturnal migration is the dominant strategy across the majority of Palearctic songbird species,^{20,21} and large-scale weather radar programs in North America have similarly focused on nocturnal movements as the established norm.^{1,22} More recently, individual-tracking studies have begun to reveal that birds may initiate flights at multiple times throughout the night, highlighting some individual variation in flight onset and duration.^{23,24} Yet, despite this wide range of methods, each comes with limitations, and none can accurately quantify diurnal and nocturnal flight activity at both the individual and species levels with detailed temporal-spatial resolution. For example, while automated radio-telemetry has been extensively used to document departure date and time, it cannot determine flight durations or landing times; similarly, acoustic monitoring fails to detect silent species, and radar monitoring is unable to distinguish between individual species.³ Visual observations of

some migratory species while they are flying during daytime may suggest diurnal migration, but these observations are often based on opportunistic encounters, and without detailed information on individual behavior, this interpretation remains speculative. As a result, while nocturnal migration is well established as the dominant strategy in many songbird groups, precisely quantifying the proportion of nocturnal versus diurnal flight at the individual and species level across a broad taxonomic and geographic range was unrealistic with earlier methods. Indeed, the literature often remains vague when classifying species as diurnal or nocturnal migrants, with many simply described as migrating during both periods (see Horton et al.⁶ and Adamík et al.²⁵). This uncertainty hampers both our theoretical understanding of diel activity rhythms and our ability to assess conservation risks, with nocturnal species being much more susceptible to light pollution, building, and wind turbine collisions.^{4–6}

Here, we used multi-sensor geolocators to obtain detailed individual behavioral tracking throughout the annual cycle. By




Figure 1. Overview of migratory flight timing across 56 small landbird species

(A) Highlights four representative individual tracks (numbered 1–4, also marked in B), showing the annual cycle locations (triangles, breeding; squares, non-breeding; circles, stopovers) and likely migration paths (gray lines), with circle size proportional to stopover duration and with the colored background reflecting

(legend continued on next page)

combining a unique set of published and unpublished tracking data, we analyzed data from 411 small- to medium-sized migratory individual landbirds to accurately quantify the proportion of migratory flights at night (Figures 1 and S1). Individuals belonged to 56 species from four continents and were from 21 families within 7 orders, and all but two species (one owl and one nightjar) were diurnal. These geolocators are ultra-light tags (averaging <1 g; see STAR Methods) that record light, barometric pressure, and, in most cases, accelerometer data but do not transmit information automatically and therefore must be retrieved to access the data.^{7,8} With data sampling intervals ranging from 5 to 30 min depending on the model, we were able to determine the precise timing of each migratory flight, defined by a clear drop in pressure data corresponding to altitude gain in flight and associated with high accelerometer measurements (of the same duration as the variation in pressure, see Nussbaumer et al.⁹ and Dhanjal-Adams et al.²⁶). We applied state-of-the-art geopositioning methods to track individual locations and then calculated civil dawn and civil dusk (i.e., first time in the morning and last time in the evening when the solar depression angle was above -6°) at different stages of each individual's annual cycle. This allowed us to determine the proportion of migratory flight that occurred during daytime versus nighttime.^{9,10}

Nocturnal migration is the dominant strategy in small- to medium-sized landbirds

Across all species, we found that, on average, 82% of all migratory flights occurred solely at nighttime (interquartile range of 79%–98%). When excluding flights involving the crossing of major ecological barriers, during which nocturnal migratory flights sometimes extended into the daytime,²⁷ this increased to 85% of migratory flights occurring at nighttime (interquartile range across species of 90%–99%). Forty-one species (out of 56; 73%) had more than 90% of their migratory flights occurring at night, confirming that nocturnal migration is the dominant strategy across a broad diversity of small landbirds (see Figure 2).

For species whose migratory timing had previously been estimated by visual observations, it was revealed that several species traditionally considered to migrate both during day and night instead conducted most of their migratory flights at night. For example, the Eurasian hoopoe *Upupa epops* and the tree pipit *Anthus trivialis* perform 91% and 71% of their migratory flights at night, respectively, whereas previous observations suggested these were instead mostly diurnal migrants (see also Briedis et al., which contradicts previous expectations for tawny pipit *Anthus campestris*,²⁸ and Vgants et al. for common starling *Sturnus vulgaris*²⁹). These findings highlight that apparent diurnal activity does not necessarily reflect the timing of migratory flight.

Yet, important exceptions to this predominance occur, with certain species showing a clear tendency to migrate during the day. In the case of the European bee-eater *Merops apiaster*, which relies on thermal updrafts to facilitate soaring flight, observing diurnal migratory activity (0.1% nocturnal flights) was expected.³⁰ However, this new approach also revealed that certain migratory flights of bee-eaters, particularly those corresponding to the crossing of the Sahara Desert, may start several hours prior to sunrise (see Figure 1A). Also, among the exceptions, hirundines have long been regarded as predominantly diurnal migrants, sometimes also making use of thermal currents. Yet, our approach reveals that the two studied hirundine species performed a substantial share of their migratory flights at night, with an average nocturnal proportion of 38%.^{31,32} Then, three alpine accentors *Prunella collaris* tracked from southern France to their breeding grounds in the Alps performed most of their migratory flights during daytime. However, this may reflect the atypical ecology of this short-distance and altitudinal migrant rather than a family-level pattern, as other Prunellidae, such as the duncock *Prunella modularis*, are regularly recorded migrating at night.^{33–35}

Finally, certain groups, such as pipits and wagtails (Motacillidae) or species like the snow bunting *Plectrophenax nivalis*, also appeared to perform more daytime flights than other species (often short flights, e.g., Richard's pipit *Anthus richardi* in Figure 1A). In the case of the snow bunting, tracked individuals bred at very high latitudes ($>78^\circ$ north) and hence conducted a substantial portion of their migratory flights during polar day. This example illustrates facultative diurnal migration under continuous daylight and reflects the same trade-off observed in high-latitude radar studies, in which preserving migratory progress takes precedence over adhering to nocturnality.³⁶

Several species in our study were represented by small sample sizes. Nevertheless, we found very little intraspecific variation, with an average interquartile range of proportion of nocturnal flights across species of 0.036 when excluding flights across ecological barriers (also 0.036 when all flights were included; Figure 3). We suspect that variation in the proportion of nocturnal flights is primarily driven by migratory route choice and the need to cross ecological barriers (see spotted flycatcher *Muscicapa striata* in Figure 1A), although species differ in their propensity to prolong flights into daytime when facing such crossings.^{27,37,38} In another example, the common starling performed its longest flights, usually during long marine crossings, mostly at night, whereas shorter flights tended to occur during the day (Figure S3).²⁹ Excluding these barrier-crossing flights tends to lower the estimated proportion of nocturnal migration and increases the estimated intraspecific variation, because

positional uncertainty. Each horizontal bar in the activity plots represents one migratory flight (flight ID, flights ordered sequentially across the annual cycle), shown relative to night periods (white bars), with a gray horizontal line separating autumn and spring migration. For individuals flying into the day (spotted flycatcher *Muscicapa striata*) or primarily during the day (European bee-eater *Merops apiaster*), two consecutive nights are included to visualize the full flight duration.

(B) Shows a phylogenetic tree alongside three bars per species: (1) the proportion of migratory flight occurring during the night (dark gray) versus day (yellow), with intra-species variation shown as interquartile ranges (black lines); (2) departure; and (3) landing times across a standardized diel cycle, with normalized nighttime centered between dusk and dawn. Alongside, we present the number of tracked individuals indicated next to each species name with a color-coded family. Dusk and dawn times are based on a solar depression angle of -6° (see STAR Methods). For species marked with an asterisk (*), flight timing was more difficult to estimate accurately due to non-stop flying or aerial feeding behavior (see STAR Methods). The proportion of migratory flights occurring at night versus during the day is estimated here, excluding barrier-crossing flights (see Figure 3A). See Figures S1 and S2, Table S1, and Data S1.

not all tracked starlings undertook such long marine crossings (Figures 3 and S3).

Most species synchronized the departure and landing of migratory flights close to civil dusk and civil dawn. Specifically, 92% and 96% of species synchronized departures and landings more closely with civil twilight than with actual sunset or sunrise, respectively, suggesting that nocturnal migrants tend to maximize travel time during the night (Figure S2).^{11,24,39} However, some species-specific exceptions exist. For instance, the three intra-tropical migrants tracked in Africa, the woodland kingfisher *Halcyon senegalensis*, the mangrove kingfisher *H. senegaloides*, and the red-capped robin-chat *Cossypha natalensis*, are exclusively nocturnal migrants, yet their departures and landings appeared less tightly synchronized with the end and start of civil twilight (Figure 1 and Rime et al.⁴⁰). A similar pattern was observed in some other species (starling, wagtails, and pipits), which also performed some of their flights during the day (Figure S3).

Phylogenetic structuring of flight timing of migratory landbirds

To describe the phylogenetic structuring of nocturnal migration and to assess whether this behavior tends to be conserved within lineages, we investigated phylogenetic signal in nocturnal flight patterns across species. Across both metrics of nocturnal flight proportions, including or excluding barrier-crossing flights, we detected a consistent phylogenetic signal, though its strength varied (Table S1). Pagel's λ was nearly identical and significant in both cases ($\lambda = 1.015$ and 1.008 , $p < 0.001$), pointing to strong phylogenetic structuring and suggesting that nocturnal migration is largely conserved within lineages. Blomberg's K^* indicated a significant and consistent signal in both cases (0.723 , $p = 0.001$ and 0.746 , $p = 0.002$), suggesting that the phylogenetic structuring of nocturnal migration is robust regardless of whether barrier-crossing flights are included. Collectively, these results show that nocturnal migration is moderately to strongly shaped by phylogeny and evolutionary history, as is the case for the tendency to evolve seasonal migration itself.^{41,42} The overall consistency across all indices and the strengthened signal in the expanded dataset reinforce the interpretation that nocturnal migration is largely conserved within lineages, though convergent evolution or behavioral plasticity under similar ecological pressures cannot be excluded.

To assess whether the detected phylogenetic signal could reflect shared ecological conditions among related species, we ran phylogenetic generalized least squares (PGLS) models incorporating breeding latitude and total migration distance as covariates (two variables *a priori* expected to influence nocturnal flight proportions, see Sivakumar et al.³⁶). Neither factor predicted nocturnal flight proportion after accounting for phylogenetic relatedness (breeding latitude: $p = 0.101$; migration distance: $p = 0.567$; Figure S4), suggesting that the phylogenetic structuring of nocturnal migration reflects true evolutionary conservatism rather than shared ecology alone.

Several hypotheses have been proposed to explain nocturnal migration, but, for the most part, they remain untested (see Alerstam¹²). A first hypothesis proposes that nighttime flights offer more favorable aerodynamic conditions, as birds avoid turbulence and strong winds.⁴³ This has also been proposed as a




Figure 2. Distribution of migratory flight timing between day and night across all pooled flights (12,542 flights) from 411 individuals, totaling 84,477 h of flight

The top panel shows the proportion of birds aloft over a normalized 24-h cycle, with 77% of the migratory flights occurring during nighttime. The middle panel illustrates normalized departure times, with shading from light pink to dark red indicating the relative percentage of departures (darker shades indicate higher concentrations). The bottom panel shows normalized landing times, with shading from light to dark blue reflecting the relative percentage of landings. Dusk and dawn times are based on a solar depression angle of -6° (see STAR Methods). All proportions within panels are rescaled between 0 and 1. See also Figure S2 and Data S1.

key driver of nocturnal migration in insects, in which radar studies have shown that nocturnal migrants achieve greater travel speeds and tighter directional control than their diurnal counterparts.^{44,45} A related hypothesis on physiological conditions proposes that nocturnal flight allows birds to avoid overheating from solar radiation exposure and from evaporative water loss because of cooler, more humid air. This is consistent with the extreme diel flight altitude changes recently documented in different species.^{27,37,46} A third hypothesis points to predator avoidance,^{47,48} and a fourth highlights the use of celestial cues after sunset and into the night for orientation,²⁴ alongside solar and magnetic cues that also provide mechanisms for orientation.^{49,50} Finally, a fifth hypothesis suggests that nocturnal migration in otherwise diurnal species evolved to maximize foraging opportunities during the day.^{51,52} This may explain why species such as hirundines, swifts, and bee-eaters, which are able to feed on the wing, distribute their migratory flights between day and night and are capable of performing rapid, uninterrupted migratory journeys in just a few days (e.g., in alpine




Figure 3. Effects of ecological barrier crossing on the mean and variation of the proportion of nocturnal migratory flight

(A) Average proportion of nocturnal migratory flights per species, with and without barrier-crossing flights. Species above the dashed line show a higher proportion of diurnal flights when crossing barriers (see STAR Methods).

(B) Intraspecific variation in nocturnal migratory activity per species, expressed as interquartile range (IQR), with and without barrier-crossing flights; species above the dashed line exhibit increased variation attributable to barrier crossing. Axes in (B) are square-root transformed for readability.

In (A) and (B), dot size is proportional to the number of tracked individuals. Only species with ≥ 3 individuals are included.

See also Figure S3 and Data S1.

swift³²). There is also some indication that Palearctic migrants adopt a more flexible timing (i.e., less tightly synchronized with civil twilight, as compared with those in North America or elsewhere) when they reach their wintering quarters south of the Sahara (see Adamík et al.⁵³). By adopting nocturnal flight strategies similar to those of intra-African migratory species, they may perform shorter, more opportunistic flights that allow them to efficiently track patchy or ephemeral food resources.^{40,54} While our data cannot directly test these hypotheses, some patterns are at least consistent with certain predictions. The near-universal synchronization of departures and landings with civil dusk and dawn is consistent with both the celestial orientation and predator avoidance hypotheses, whereas the exceptions (e.g., aerial feeders) align more naturally with the foraging hypothesis. Future studies combining multi-sensor tracking with environmental data such as wind, temperature, and predator activity would allow more direct tests of these competing explanations.

Conclusion

Our study highlights the unique value and spatiotemporal resolution of multi-sensor geolocators as the most suitable method currently available to quantify the proportion of nocturnal versus diurnal migration in small- to medium-sized landbirds and also to investigate their behavior en route. This approach, however, comes with inherent limitations because it relies on species that have sufficient site fidelity to enable archival tag recovery in subsequent years. As a result, our dataset is subject to taxonomic and geographic biases, with certain groups (e.g.,

Passerellidae and Tyrannidae) remaining underrepresented (Figure 4). Among the groups most notably absent from our dataset are the Fringillidae, of which several members (e.g., crossbills and siskins) are presumed, from visual observations and ringing studies, to undertake substantial daytime migration. Including such species in future analyses would likely reveal a broader continuum of diurnal and nocturnal strategies than captured here and would refine estimates of the overall prevalence of nocturnal migration across landbirds. Expanding taxonomic coverage to currently underrepresented groups would also allow testing whether the phylogenetic conservatism detected here holds across a broader avian tree or whether certain lineages have independently evolved flexible diel strategies. Furthermore, combining multi-sensor tracking with concurrent environmental measurements (e.g., wind, temperature, and predator activity) would enable direct evaluation of the competing mechanistic hypotheses for nocturnal migration that our descriptive framework cannot resolve. By providing a synthesis of both published and unpublished datasets, we hope this work will encourage new collaborations, promote wider use of multi-sensor tracking technology, and draw attention to the current taxonomic and geographic gaps.

RESOURCE AVAILABILITY

Lead contact

Requests for further information and resources should be directed to and will be fulfilled by the lead contact, Paul Dufour (paul.dufour80@gmail.com).




Figure 4. Phylogenetic distribution of multi-sensor tracking data, nocturnality, and diurnality across migratory landbird families

The phylogenetic tree includes all migratory landbird families. For each family, the outer circle indicates whether multi-sensor tracking data are available and, if so, the average proportion of nocturnal migratory flights across tracked species within that family, represented by a color ranging from yellow (>50% diurnal) to dark gray (>50% nocturnal). White circles indicate families for which no multi-sensor tracking data are currently available. Circle size is proportional to the total number of known migratory species within each family. Migration strategies were taken from Dufour et al.⁵⁵ Inner colored dots indicate the geographic distribution of each family (pink, Old World and Oceania; blue, New World; green: both). Family names and their corresponding numbers are listed to the right of the figure. See also [Data S1](#).

Materials availability

This study did not generate new unique reagents.

Data and code availability

- All data necessary to reproduce this publication are available in [Data S1](#). It provides information about the estimated proportion of nocturnal flights for each individual.
- The study does not report any original code.
- Any additional information required to reanalyze the data reported in this paper is available from the [lead contact](#) upon request.

ACKNOWLEDGMENTS

We thank Anna Anashina, Miroslav Bažant, Johanna Beam, Václav Beran, Oleg Bourski, Vojtěch Brlík, Olivia Butler, Wenjia Chen, Ming Cheng, Paul Coiffard, Yannig Coulomb, Rose Delacroix, Julie Deschamps, Dimitar Dimitrov, Ivo Dinsbergs, Dule, Tamara Emmenegger, Ignacio C. Fernández Calvo, Christophe de Franceschi, Yayue Gao, Sophie de Grissac, Felipe González Sánchez, Jan Grünwald, Steffen Hahn, Dennis Hasselquist, Javier Hernández Cabello, Rob Husbands, Mihaela Ilieva, Ilya Ishchenko, Karina Ivanova, Māris Jaunzemis, Valts Jaunzemis, Bastien Jeannin, Shengjing Jiao, Oskars Keiřs, Dmitry Kishkinev, Jaroslav Koleček, Anton Kristin, Thibaut Lacombe, Fumin Lei, Dmitry Loke, Zhuo Li, Danmeng Liu, Mads Moore, Martin Marinov, Robin Monchatre, Lykke Pedersen, Katie Percy, Robbie Phillips, Strahil Peev, Milica Požgayová, Ance Priedniece, Fanny Rey, Jeff Roth, Benjamin Rousseau, Ekaterina Schemeleva, Christoffer Sjöholm, Katherine R.S. Snell, Gang Song, Kateřina Sosnovcová, Christian Stolz, Michal Šulc, Stephan Tillo, Viesturs Vgants, Xinyuan Wang, Elza Zacmane, Pavel Zehindjiev, Guoming Zhang,

and Yu Zhang for their help in deploying and retrieving the loggers; James Fox, Erich Bächler, Neringa Znakovaitė, Arne Andersson, and Johan Bäckman for their help in extracting the data and in developing the geolocators and software; and Mathieu Gravey for his help developing the GeoPressureR package for the analysis of this study. A.B.R. and G.P.d.S. were funded via the “Santander Capital Natural” project and SEO/Birdlife. P.C.-L. is funded by a global Marie Skłodowska-Curie Actions Fellowship (HORIZON-MSCA-2023-PF-01-101150591). Most birds tagged in France, and some in the UK, were part of the Migration and Migratlane research projects, which were funded by the French minister of Ecology and the Office Français de la Biodiversité (OFB). The development of multi-sensor loggers at Lund University received support from the Centre for Animal Movement Research (CAnMove), financed by a Linnaeus Grant (349-2007-8690) from the Swedish Research Council and Lund University. M. Briedis received funding from the Latvian Council of Science (project no. lzp-2023/1-0233). P.P. was supported by the Institutional Research Plan (RVO: 68081766). P.A. was supported by the Palacký University grant scheme (IGA_PrF_2025_26). N.C. received funding from Smithsonian (0100024) and the USFWS (F24AP00033-00). M. Liedvogel received funding from the DFG (project Nav05 within SFB 1372-Magnetoreception and Navigation in Vertebrates, project no 395940726). S. Sjöberg received funding from the European Research Council (ERC starting grant no. 101078349) and the Swedish Research Council (starting grant no. 2021-05654). P.D.-V. was supported by the Fondation Biotope. A.H. received funding from the Swedish Research Council (2020-03707) and the Knut and Alice Wallenberg Foundation (KAW 2020.0096). S.A. received funding from the Swedish Research Council (2016-05342 and 2022-03755) and the Royal Physiographical Society in Lund. E.I.J. received funding from the Coypu Foundation. S. Szarmach was awarded research grants from the American Ornithological Society, the Animal Behavior Society, and the Wilson Ornithological Society, as well as an

Achievement Graduate Fellowship and a Department of Biology and Huck Institute of Life Sciences grant at Pennsylvania State University.

AUTHOR CONTRIBUTIONS

P.D. designed the paper with R.N., B.H., and M. Briedis. P.D. performed analyses, visualizations, and wrote the first version of the paper. P.D., R.N., J.B.W., Y.R., P.M.-T., M. Briedis, E.J., V.C.-L., P.C.-L., G.R., T.Z., P.D.-V., S. Szarmach, D.K., and K.J.G. analyzed tracking data. All authors were involved in the data collection in the field. All authors contributed to the revisions.

DECLARATION OF INTERESTS

The authors declare no competing interests.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS
- METHOD DETAILS
 - Geolocators and species data
 - Estimations of nocturnal and diurnal flight times
 - Cases of aerial feeders and non-stop flying species
 - Ecological barrier crossing flights
- QUANTIFICATION AND STATISTICAL ANALYSIS
 - Phylogenetic modelling

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.cub.2026.06.030>.

Received: October 11, 2025

Revised: April 16, 2026

Accepted: June 10, 2026

Published: July 8, 2026

REFERENCES

1. Dokter, A.M., Farnsworth, A., Fink, D., Ruiz-Gutierrez, V., Hochachka, W.M., La Sorte, F.A., Robinson, O.J., Rosenberg, K.V., and Kelling, S. (2018). Seasonal abundance and survival of North America's migratory avifauna determined by weather radar. *Nat. Ecol. Evol.* 2, 1603–1609. <https://doi.org/10.1038/s41559-018-0666-4>.
2. Mukhin, A., Grinkevich, V., and Helm, B. (2009). Under Cover of Darkness: Nocturnal Life of Diurnal Birds. *J. Biol. Rhythms* 24, 225–231. <https://doi.org/10.1177/0748730409335349>.
3. Hüppop, O., Ciach, M., Diehl, R., Reynolds, D.R., Stepanian, P.M., and Menz, M.H.M. (2019). Perspectives and challenges for the use of radar in biological conservation. *Ecography* 42, 912–930. <https://doi.org/10.1111/ecog.04063>.
4. Burt, C.S., Kelly, J.F., Trankina, G.E., Silva, C.L., Khalighifar, A., Jenkins-Smith, H.C., Fox, A.S., Fristrup, K.M., and Horton, K.G. (2023). The effects of light pollution on migratory animal behavior. *Trends Ecol. Evol.* 38, 355–368. <https://doi.org/10.1016/j.tree.2022.12.006>.
5. Van Doren, B.M., Horton, K.G., Dokter, A.M., Klinck, H., Elbin, S.B., and Farnsworth, A. (2017). High-intensity urban light installation dramatically alters nocturnal bird migration. *Proc. Natl. Acad. Sci. USA* 114, 11175–11180. <https://doi.org/10.1073/pnas.1708574114>.
6. Horton, K.G., Nilsson, C., Van Doren, B.M., La Sorte, F.A., Dokter, A.M., and Farnsworth, A. (2019). Bright lights in the big cities: migratory birds' exposure to artificial light. *Front. Ecol. Environ.* 17, 209–214. <https://doi.org/10.1002/fee.2029>.
7. Liechti, F., Bauer, S., Dhanjal-Adams, K.L., Emmenegger, T., Zehndtjiev, P., and Hahn, S. (2018). Miniaturized multi-sensor loggers provide new insight into year-round flight behaviour of small trans-Saharan avian migrants. *Mov. Ecol.* 6, 19. <https://doi.org/10.1186/s40462-018-0137-1>.
8. Sjöberg, S., Pedersen, L., Malmiga, G., Alerstam, T., Hansson, B., Hasselquist, D., Thorup, K., Tøttrup, A.P., Andersson, A., and Bäckman, J. (2018). Barometer logging reveals new dimensions of individual songbird migration. *J. Avian Biol.* 49, e01821. <https://doi.org/10.1111/jav.01821>.
9. Nussbaumer, R., Gravey, M., Briedis, M., and Liechti, F. (2023). Global positioning with animal-borne pressure sensors. *Methods Ecol. Evol.* 14, 1104–1117. <https://doi.org/10.1111/2041-210X.14043>.
10. Nussbaumer, R., Gravey, M., Briedis, M., Liechti, F., and Sheldon, D. (2023). Reconstructing bird trajectories from pressure and wind data using a highly optimized hidden Markov model. *Methods Ecol. Evol.* 14, 1118–1129. <https://doi.org/10.1111/2041-210X.14082>.
11. Moore, F.R. (1987). Sunset and the orientation behavior of migrating birds. *Biol. Rev.* 62, 65–86. <https://doi.org/10.1111/j.1469-185X.1987.tb00626.x>.
12. Alerstam, T. (2009). Flight by night or day? Optimal daily timing of bird migration. *J. Theor. Biol.* 258, 530–536. <https://doi.org/10.1016/j.jtbi.2009.01.020>.
13. Alerstam, T. (1990). *Bird Migration* (Cambridge University Press).
14. Helm, B., Visser, M.E., Schwartz, W., Kronfeld-Schor, N., Gerkema, M., Piersma, T., and Bloch, G. (2017). Two sides of a coin: ecological and chronobiological perspectives of timing in the wild. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 372, 20160246. <https://doi.org/10.1098/rstb.2016.0246>.
15. Gilbert, N.A., McGinn, K.A., Nunes, L.A., Shipley, A.A., Bernath-Plaisted, J., Clare, J.D.J., Murphy, P.W., Keyser, S.R., Thompson, K.L., Maresch Nelson, S.B., et al. (2023). Daily activity timing in the Anthropocene. *Trends Ecol. Evol.* 38, 324–336. <https://doi.org/10.1016/j.tree.2022.10.008>.
16. Van Doren, B.M., Løstam, V., Cramer, A., Salamon, J., Dokter, A., Kelling, S., Bello, J.P., and Farnsworth, A. (2023). Automated acoustic monitoring captures timing and intensity of bird migration. *J. Appl. Ecol.* 60, 433–444. <https://doi.org/10.1111/1365-2664.14342>.
17. Gwinner, E. (1986). *Circannual Rhythms* (Springer). <https://doi.org/10.1007/978-3-642-82870-6>.
18. Horton, K.G., La Sorte, F.A., Sheldon, D., Lin, T.-Y., Winner, K., Bernstein, G., Maji, S., Hochachka, W.M., and Farnsworth, A. (2020). Phenology of nocturnal avian migration has shifted at the continental scale. *Nat. Clim. Chang.* 10, 63–68. <https://doi.org/10.1038/s41558-019-0648-9>.
19. Johnson, T.B., Gochfeld, D., Roach, M.C., Apte, G.A., Gallardy, R., Wilson, S.C., Marden, A., Irons, D.J., Farnsworth, A., and Rutt, C.L. (2026). Seeing in the dark: Using thermal imaging to directly observe nocturnal migration. *Ornithology* 143, 1–9. <https://doi.org/10.1093/ornithology/ukaf074>.
20. Schmaljohann, H., Liechti, F., and Bruderer, B. (2007). Songbird migration across the Sahara: the non-stop hypothesis rejected! *Proc. Biol. Sci.* 274, 735–739. <https://doi.org/10.1098/rspb.2006.0011>.
21. Bruderer, B., and Liechti, F. (1998). Flight Behaviour of Nocturnally Migrating Birds in Coastal Areas: Crossing or Coasting. *J. Avian Biol.* 29, 499. <https://doi.org/10.2307/3677169>.
22. Van Doren, B.M., and Horton, K.G. (2018). A continental system for forecasting bird migration. *Science* 361, 1115–1118. <https://doi.org/10.1126/science.aat7526>.
23. Müller, F., Taylor, P.D., Sjöberg, S., Muheim, R., Tsvey, A., Mackenzie, S.A., and Schmaljohann, H. (2016). Towards a conceptual framework for explaining variation in nocturnal departure time of songbird migrants. *Mov. Ecol.* 4, 24. <https://doi.org/10.1186/s40462-016-0089-2>.
24. Cooper, N.W., Dossman, B.C., Berrigan, L.E., Brown, J.M., Brunner, A.R., Chmura, H.E., Cormier, D.A., Bégin-Marchand, C., Rodewald, A.D., Taylor, P.D., et al. (2023). Songbirds initiate migratory flights synchronously relative to civil dusk. *Mov. Ecol.* 11, 24. <https://doi.org/10.1186/s40462-023-00382-5>.

25. Adamik, P., Emmenegger, T., Briedis, M., Gustafsson, L., Henshaw, I., Krist, M., Laaksonen, T., Liechti, F., Procházka, P., Salewski, V., et al. (2016). Barrier crossing in small avian migrants: individual tracking reveals prolonged nocturnal flights into the day as a common migratory strategy. *Sci. Rep.* 6, 21560. <https://doi.org/10.1038/srep21560>.
26. Dhanjal-Adams, K.L., Willener, A.S.T., and Liechti, F. (2022). pamlr: a toolbox for analysing animal behaviour using pressure, acceleration, temperature, magnetic or light data in R. *J. Anim. Ecol.* 91, 1345–1360. <https://doi.org/10.1111/1365-2656.13695>.
27. Dufour, P., Nussbaumer, R., Briedis, M., Bocher, P., Conway, G., Coulomb, Y., Delacroix, R., Dagonet, T., de Franceschi, C., de Grissac, S., et al. (2026). Ecological barrier crossing strategies in small migratory birds depend on wing morphology and plumage color. *iScience* 29, 114466. <https://doi.org/10.1016/j.isci.2025.114466>.
28. Briedis, M., Beran, V., Adamik, P., and Hahn, S. (2020). Integrating light-level geolocation with activity tracking reveals unexpected nocturnal migration patterns of the tawny pipit. *J. Avian Biol.* 51, e02546. <https://doi.org/10.1111/jav.02546>.
29. Vgants, V., Keiss, O., Dinsbergs, I., Jaunzemis, V., Zacmane, E.M., Priedniece, A., and Briedis, M. (2023). Migration strategies, performance and annual activity budget in a short-distance migrant, the Common Starling *Sturnus vulgaris*. *J. Avian Biol.* 2023, e03080. <https://doi.org/10.1111/jav.03080>.
30. Sapir, N., Wikelski, M., Avissar, R., and Nathan, R. (2011). Timing and flight mode of departure in migrating European bee-eaters in relation to multi-scale meteorological processes. *Behav. Ecol. Sociobiol.* 65, 1353–1365. <https://doi.org/10.1007/s00265-011-1146-x>.
31. Karlsson, H., Henningsson, P., Bäckman, J., Hedenström, A., and Ålerstam, T. (2010). Compensation for wind drift by migrating swifts. *Anim. Behav.* 80, 399–404. <https://doi.org/10.1016/j.anbehav.2010.05.023>.
32. Liechti, F., Witvliet, W., Weber, R., and Bächler, E. (2013). First evidence of a 200-day non-stop flight in a bird. *Nat. Commun.* 4, 2554. <https://doi.org/10.1038/ncomms3554>.
33. Heiss, M. (2018). Nocturnal bird migration at Besh Barnag bottleneck in Azerbaijan as revealed by means of acoustic monitoring. *Bird Conserv. Int.* 28, 593–605. <https://doi.org/10.1017/S0959270917000454>.
34. Schwabl, H., Gwinner, E., Benvenuti, S., and Ioalè, P. (1991). Exposure of Dunnocks (*Prunella modularis*) to their Previous Wintering Site Modifies Autumnal Activity Pattern: Evidence for Site Recognition? *Ethology* 88, 35–45. <https://doi.org/10.1111/j.1439-0310.1991.tb00261.x>.
35. Klinner, T., Karwinkel, T., Packmor, F., and Schmaljohann, H. (2025). Stopover departure decisions in spring: pre-Saharan migrants stay longer and are more selective for favourable wind than trans-Saharan migrants. *Mov. Ecol.* 13, 64. <https://doi.org/10.1186/s40462-025-00575-0>.
36. Sivakumar, A.H., Sheldon, D., and Horton, K.G. (2025). High-latitude night shortening forces birds to accept increasingly diurnal migration to maintain flight duration. *J. Avian Biol.* 2025, e03405. <https://doi.org/10.1002/jav.03405>.
37. Sjöberg, S., Andersson, A., Bäckman, J., Hansson, B., Malmiga, G., Tarka, M., Hasselquist, D., Lindström, Å., and Ålerstam, T. (2023). Solar heating may explain extreme diel flight altitude changes in migrating birds. *Curr. Biol.* 33, 4232–4237.e2. <https://doi.org/10.1016/j.cub.2023.08.035>.
38. Sokolovskis, K., Caballero-Lopez, V., Åkesson, S., Lundberg, M., Willemoes, M., Zhao, T., and Bensch, S. (2023). Diurnal migration patterns in willow warblers differ between the western and eastern flyways. *Mov. Ecol.* 11, 58. <https://doi.org/10.1186/s40462-023-00425-x>.
39. Åkesson, S., Ålerstam, T., and Hedenström, A. (1996). Flight Initiation of Nocturnal Passerine Migrants in Relation to Celestial Orientation Conditions at Twilight. *J. Avian Biol.* 27, 95. <https://doi.org/10.2307/3677138>.
40. Rime, Y., Osinubi, S.T., Liechti, F., Helm, B., and Nussbaumer, R. (2025). Small intra-tropical long-distance migratory birds track rainy seasons across hemispheres. *Proc. Biol. Sci.* 292, 20242633. <https://doi.org/10.1098/rspb.2024.2633>.
41. Winger, B.M., Barker, F.K., and Ree, R.H. (2014). Temperate origins of long-distance seasonal migration in New World songbirds. *Proc. Natl. Acad. Sci. USA* 111, 12115–12120. <https://doi.org/10.1073/pnas.1405000111>.
42. Dufour, P., Descamps, S., Chantepie, S., Renaud, J., Guéguen, M., Schiffrers, K., Thuiller, W., and Lavergne, S. (2020). Reconstructing the geographic and climatic origins of long-distance bird migrations. *J. Biogeogr.* 47, 155–166. <https://doi.org/10.1111/jbi.13700>.
43. Kerlinger, P., and Moore, F.R. (1989). Atmospheric Structure and Avian Migration. In *Current Ornithology*, D.M. Power, ed. (Springer), pp. 109–142. https://doi.org/10.1007/978-1-4757-9918-7_3.
44. Chapman, J.W., Drake, V.A., and Reynolds, D.R. (2011). Recent Insights from Radar Studies of Insect Flight. *Annu. Rev. Entomol.* 56, 337–356. <https://doi.org/10.1146/annurev-ento-120709-144820>.
45. Hu, G., Lim, K.S., Reynolds, D.R., Reynolds, A.M., and Chapman, J.W. (2016). Wind-Related Orientation Patterns in Diurnal, Crepuscular and Nocturnal High-Altitude Insect Migrants. *Front. Behav. Neurosci.* 10, 32. <https://doi.org/10.3389/fnbeh.2016.00032>.
46. Lindström, Å., Ålerstam, T., Andersson, A., Bäckman, J., Bählenberg, P., Bom, R., Ekblom, R., Klaassen, R.H.G., Korniluk, M., Sjöberg, S., et al. (2021). Extreme altitude changes between night and day during marathon flights of great snipes. *Curr. Biol.* 31, 3433–3439.e3. <https://doi.org/10.1016/j.cub.2021.05.047>.
47. Xirouchakis, S.M., and Panuccio, M. (2019). Hunting Altitude of Eleonora's Falcon (*Falco eleonora*) Over a Breeding Colony. *J. Raptor Res.* 53, 56. <https://doi.org/10.3356/JRR-17-94>.
48. Boom, M.P., Yu, H., Bom, R.A., Hegemann, A., Lindström, Å., Nolet, B.A., and Lameris, T.K. (2024). Migrating shorebird killed by raptor at 3000 m above ground as revealed by high-resolution tracking. *Ecology* 105, e4437. <https://doi.org/10.1002/ecy.4437>.
49. Åkesson, S., and Helm, B. (2020). Endogenous Programs and Flexibility in Bird Migration. *Front. Ecol. Evol.* 8, 78. <https://doi.org/10.3389/fevo.2020.00078>.
50. Prinz, D., Heim, R.J., Meinken, M., Niemann, N., Temme, L., Esther, A., and Heim, W. (2025). Lunar cycle and moonlight intensity influence nocturnal migration patterns in a small songbird. *Sci. Rep.* 15, 19944. <https://doi.org/10.1038/s41598-025-04270-3>.
51. Lank, D.B. (1989). Why Fly by Night? Inferences from Tidally-Induced Migratory Departures of Sandpipers. *J. Field Ornithol.* 60, 154–161.
52. Hedenström, A. (2025). Integrating flight mechanics, energetics and migration ecology in vertebrates. *J. Exp. Biol.* 228, JEB248123. <https://doi.org/10.1242/jeb.248123>.
53. Adamik, P., Wong, J.B., Hahn, S., and Kristin, A. (2024). Non-breeding sites, loop migration and flight activity patterns over the annual cycle in the Lesser Grey Shrike *Lanius minor* from a north-western edge of its range. *J. Ornithol.* 165, 247–256. <https://doi.org/10.1007/s10336-023-02102-7>.
54. Thorup, K., Tøttrup, A.P., Willemoes, M., Klaassen, R.H.G., Strandberg, R., Vega, M.L., Dasari, H.P., Araújo, M.B., Wikelski, M., and Rahbek, C. (2017). Resource tracking within and across continents in long-distance bird migrants. *Sci. Adv.* 3, e1601360. <https://doi.org/10.1126/sciadv.1601360>.
55. Dufour, P., Sayol, F., Cooke, R., Blackburn, T.M., Gallien, L., Griesser, M., Steinbauer, M.J., and Faurby, S. (2024). The importance of migratory drop-off for island colonization in birds. *Proc. Biol. Sci.* 291, 20232926. <https://doi.org/10.1098/rspb.2023.2926>.
56. Miller, E., McTavish, E., and Sanchez-Reyes, L. (2025). clooti: Fetch and Explore the Cornell Lab of Ornithology Open Tree of Life Avian Phylogeny. <https://github.com/eliottmiller/clooti>.
57. Pinheiro, J., and Bates, D. (1999). nlme: Linear and Nonlinear Mixed Effects Models. <https://doi.org/10.32614/CRAN.package.nlme>.
58. Keck, F., Rimet, F., Bouchez, A., and Franc, A. (2016). phylosignal: an R package to measure, test, and explore the phylogenetic signal. *Ecol. Evol.* 6, 2774–2780. <https://doi.org/10.1002/ecs3.2051>.

59. Hijmans, R., Williams, E., and Vennes, C. (2015). Geosphere: spherical trigonometry. R package. <https://cran.r-project.org/web/packages/geosphere/index.html>.
60. Bäckman, J., Andersson, A., Alerstam, T., Pedersen, L., Sjöberg, S., Thorup, K., and Tøttrup, A.P. (2017). Activity and migratory flights of individual free-flying songbirds throughout the annual cycle: method and first case study. *J. Avian Biol.* 48, 309–319. <https://doi.org/10.1111/jav.01068>.
61. Rhyne, G.S., Stouffer, P.C., Briedis, M., and Nussbaumer, R. (2024). Barometric geolocators can reveal unprecedented details about the migratory ecology of small birds. *Ornithology* 141, ukae010. <https://doi.org/10.1093/ornithology/ukae010>.
62. Lisovski, S., and Hahn, S. (2012). Geolight - processing and analysing light-based geolocator data in R. *Methods Ecol. Evol.* 3, 1055–1059. <https://doi.org/10.1111/j.2041-210X.2012.00248.x>.
63. Lisovski, S., Hewson, C.M., Klaassen, R.H.G., Korner-Nievergelt, F., Kristensen, M.W., and Hahn, S. (2012). Geolocation by light: accuracy and precision affected by environmental factors. *Methods Ecol. Evol.* 3, 603–612. <https://doi.org/10.1111/j.2041-210X.2012.00185.x>.
64. Lisovski, S., Bauer, S., Briedis, M., Davidson, S.C., Dhanjal-Adams, K.L., Hallworth, M.T., Karagicheva, J., Meier, C.M., Merkel, B., Ouwehand, J., et al. (2020). Light-level geolocator analyses: A user's guide. *J. Anim. Ecol.* 89, 221–236. <https://doi.org/10.1111/1365-2656.13036>.
65. Hedenström, A., Norevik, G., Warfvinge, K., Andersson, A., Bäckman, J., and Åkesson, S. (2016). Annual 10-Month Aerial Life Phase in the Common Swift *Apus apus*. *Curr. Biol.* 26, 3066–3070. <https://doi.org/10.1016/j.cub.2016.09.014>.
66. Loonstra, A.H.J., Verhoeven, M.A., Senner, N.R., Both, C., and Piersma, T. (2019). Adverse wind conditions during northward Sahara crossings increase the in-flight mortality of Black-tailed Godwits. *Ecol. Lett.* 22, 2060–2066. <https://doi.org/10.1111/ele.13387>.
67. Guo, F., Buler, J.J., Dokter, A.M., Horton, K.G., Cohen, E.B., Sheldon, D., Smolinsky, J.A., and Wilcove, D.S. (2025). Assessing the Corn Belt as an anthropogenic barrier to migrating landbirds in the United States. *Conserv. Biol.* 39, e70070. <https://doi.org/10.1111/cobi.70070>.
68. Brito, J.C., Tarroso, P., Vale, C.G., Martínez-Freiria, F., Boratyński, Z., Campos, J.C., Ferreira, S., Godinho, R., Gonçalves, D.V., Leite, J.V., et al. (2016). Conservation Biogeography of the Sahara-Sahel: additional protected areas are needed to secure unique biodiversity. *Divers. Distrib.* 22, 371–384. <https://doi.org/10.1111/ddi.12416>.
69. McTavish, E.J., Gerbracht, J.A., Holder, M.T., Iliff, M.J., Lepage, D., Rasmussen, P.C., Redelings, B.D., Sánchez Reyes, L.L., and Miller, E.T. (2025). A complete and dynamic tree of birds. *Proc. Natl. Acad. Sci. USA* 122, e2409658122. <https://doi.org/10.1073/pnas.2409658122>.
70. Münkemüller, T., Lavergne, S., Bzeznik, B., Dray, S., Jombart, T., Schiffrers, K., and Thuiller, W. (2012). How to measure and test phylogenetic signal. *Methods Ecol. Evol.* 3, 743–756. <https://doi.org/10.1111/j.2041-210X.2012.00196.x>.
71. Blomberg, S.P., and Garland, T. (2002). Tempo and mode in evolution: phylogenetic inertia, adaptation and comparative methods. *J. Evol. Biol.* 15, 899–910. <https://doi.org/10.1046/j.1420-9101.2002.00472.x>.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Deposited data		
Estimated proportions of nocturnal migratory flights per individuals, with or without barrier crossing flights	This paper, Data S1	DOI of the paper
Software and algorithms		
Rstudio	RStudio Team https://www.rstudio.com/	Rstudio 4.4.1
R package GeoPressureR	Nussbaumer et al. ^{9,10}	https://github.com/Rafnuss/GeoPressureR
R package clootl	Miller et al. ⁵⁶	https://cran.r-project.org/web/packages/clootl/index.html
R package nlme	Pinheiro and Bates ⁵⁷	https://cran.r-project.org/web/packages/nlme/index.html
R package phylosignal	Keck et al. ⁵⁸	https://cran.r-project.org/web/packages/phylosignal/index.html
R package geosphere	Hijmans et al. ⁵⁹	https://cran.r-project.org/web/packages/geosphere/index.html
Other		
Multi-sensor geolocators	Swiss Ornithological Institute	GDL3-PAM 1.2-2g
Multi-sensor geolocators	Migrate Technology	BARP30Z11-7-DIP, DARK30Z11-DIP, CARP30Z11-7-DIP, CARP30B1-7-DIP 0.3-0.6g
Multi-sensor geolocators	Lund University	CAnMove 1g

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

All birds were captured and tagged for the purposes of this study under permits issued by museums or the relevant local authorities in each participating country or region involved.

METHOD DETAILS

Geolocators and species data

We collected both published and unpublished multi-sensor tracking data from 411 individuals of 56 different species, originating from 83 populations across 25 countries spanning four continents (see [Figure S1](#); [Table S2](#)). We collected varying numbers of loggers per species, with a minimum of one for five species ([Table S2](#)) and a maximum of 35 for the great reed warbler *Acrocephalus arundinaceus* (7.4 loggers per species on average). Note that in some cases of partial migratory species, such as the meadow pipit *Anthus pratensis*, seven loggers were retrieved from the same population, but only one bird migrated.

Multi-sensor loggers refer to small devices that can include sensors such as light, accelerometer, and pressure. After deployment, the same individuals must be recaptured in subsequent years to recover the tags and access the data. Except for alpine accentor in France, Richard's pipit in Spain and red-capped robin-chat and mangrove kingfisher in Kenya, which were tagged on their wintering grounds, all individuals were tagged on their breeding grounds. Therefore, all tagged birds were experienced individuals that had completed at least one-way migration before tagging.

The data were collected using different types of multi-sensor loggers, which always included at least a pressure sensor, with varying recording intervals ranging from 5 to 30 minutes. These loggers were manufactured by the Swiss Ornithological Institute⁷ (model GDL3-PAM, 1.2–2.0 g with harness), the Lund University^{8,60} (<1.2 g with harness), or Migrate Technology (models BARP30Z11-7-DIP, DARK30Z11-DIP, CARP30Z11-7-DIP, CARP30B1-7-DIP, with respective weights of 0.35 and <0.55 g including harness). Most models featured at least light, accelerometer, and pressure sensors, though some (e.g., DARK30Z11-DIP) only included pressure sensors. When present, light data were always recorded at 5-minute intervals. Pressure and activity data were recorded at intervals ranging from 5 to 30 minutes.

The species tagged ranged in weight from 7 g (the smallest, common chiffchaff *Phylloscopus collybita*) to 150 g (the heaviest, European roller *Coracias garrulus*). The devices never exceeded 5% of the body mass of the tagged birds. All devices were mounted

using leg-loop harnesses made of either stretchy elastic thread or tressed UV-proof thread, with diameters between 0.7 and 1.0 mm depending on the species' size. Birds were captured and tagged following strict ethical guidelines and under various licenses issued by national or regional authorities. A list of reference programs or project codes for unpublished data is provided in the [acknowledgments](#).

Estimations of nocturnal and diurnal flight times

To determine what fraction of each migratory flight occurred during the night versus the day, we first modeled the trajectory of each track following the approach presented in Nussbaumer et al.^{9,10} using the *GeoPressureR* R package (version 3.2). This package is based on geolocation estimation of stopover sites (as well as wintering or breeding areas) throughout the annual cycle, using either pressure data alone or a combination of pressure and light data. In general, we followed all the steps presented in the online tutorial: <https://raphaelnussbaumer.com/GeoPressureManual/>.

First, we separated stationary periods and migratory flights by manually labelling the geolocators' pressure and/or activity (i.e., accelerometer) measurements. Stationary periods were characterized by a limited variation in consecutive pressure measurements indicating an absence of change in altitude as well as by consistently low activity measurements. Migratory flights typically displayed a clear drop in atmospheric pressure, corresponding to in-flight altitude gain, and consistently high activity measurements (which forms a high activity plateau of the same duration as the variation in pressure). If both pressure and activity data are available, labeling is done using activity data; otherwise, it is done using pressure data. To enable comparison between all individuals, especially those with different data recording intervals, we only labeled flights that lasted at least 40 minutes. See <https://raphaelnussbaumer.com/GeoPressureManual/labelling-tracks.html> and⁶¹ for examples.

Second, we built separate probability maps based on atmospheric pressure and light intensity data (sunrise and sunset times) to estimate the position of each bird during each stationary period. For the pressure-based maps, the time series of the geolocator pressure measurements during stationary periods were matched with the one-hour atmospheric pressure at surface level from ERA5 re-analysis dataset (spatial resolution: $0.1 \times 0.1^\circ$) to produce a likelihood map of the geolocator's position.⁹ The resulting likelihood map included the information of both the temporal variation of pressure and the absolute values of pressure corresponding to the altitudinal range within each grid cell. Stationary periods identified through pressure and/or activity measurement labeling were directly used to estimate the probability maps calculated with light intensity data. For those, we calculated likelihood estimates using an "in-habitat" calibration from the equipment and retrieval periods (as described in Lisovski and Hahn⁶² and Lisovski et al.⁶³) fitting the distribution of zenith angle with a kernel density estimation. Note that we did not use light intensity data (but only atmospheric pressure) for nocturnal species (like Eurasian scops-owl *Otus scops*) or species spending most of daytime under cover (like siberian ruby-throat *Calliope calliope*; see later for details about all individuals).

Finally, we constructed the trajectories of each bird using a Hidden Markov Model as presented in Nussbaumer et al.¹⁰ The observation model consisted of the likelihood maps generated from pressure and light data (or only pressure data). The movement model used the information of flight duration derived from the labelling and a parametric equation defined as the cubic root of the mechanical power required for the average airspeed computed for a transition, accounting for the different species size and shape. Using this model, we generated the most likely migration trajectory of the birds.

From the most likely trajectory, we then used the *path2twilight* function in the *GeoPressureR* R package to calculate the theoretical twilight times at each location for specified dates. Since civil twilight is a commonly used celestial cue (see Cooper et al.²⁴), we estimated twilight times using a solar depression angle of -6° . We also calculated twilight times using a solar depression angle of 0° , corresponding to sunrise and sunset times, to assess whether departure and landing events align more closely with civil twilight than with sunrise or sunset. Once we had the start and end of civil twilight estimated at each stopover location, we used the *stap2-flight* function to convert the stationary period data frame into a flight data frame where flights are computed as the difference between the end of a stationary period to the start of the next one. For each flight, using the civil twilight information, we then estimated the duration of the flight that occurred during the night versus the day.

Note that we favored this method of using estimation of geo-positioning over direct calculation from raw light data to estimate civil twilight times because it provides consistency across all individuals, especially since some species or types of loggers either do not record light data or produce data of insufficient quality. Although there is inherent uncertainty in geolocation, we expect the biases to be relatively small, given that variations in civil twilight times are minimal across relatively large geographic distances.

Cases of aerial feeders and non-stop flying species

Unlike species that undertake long flight phases only during migration, aerial insectivores, such as swallows, swifts, and bee-eaters, spend most of the day flying while foraging. This makes it harder to identify migratory flights from foraging flights using the pattern described in other species: a drop in pressure data combined with a sustained plateau of high activity.

Using pressure and accelerometer data, we found that most species (except swifts) typically perch at ground level for most nights. We therefore defined migratory flights as pronounced daytime activity periods (high accelerometer data) occurring between two nights that differed in mean atmospheric pressure by at least 3 hPa. Following this labeling approach and combining it with light-level data, we used *GeoPressureR* to estimate the most likely migration trajectory. We iteratively refined the labeling process until the resulting trajectory appeared coherent and did not suggest numerous back-and-forth movements, which would imply that the total duration of labeled flight was overestimated relative to the time required for the migration (see Nussbaumer et al.¹⁰).

For the only non-stop flyer in our dataset, the Alpine Swift, we first estimated the migration route, including main wintering and major stopover sites, from light data using GeoLight following Lisovski et al.⁶⁴ We then identified migration periods from shifts in sunrise and sunset, isolated these intervals in the pressure and accelerometer data, and manually labeled segments consistent with migratory behavior. These segments were marked by prolonged high-activity plateaus in the accelerometer data, probably induced by increased flapping flights, which were not observed outside migration periods. The trajectory was then interpolated to match flight events, and local sunrise/sunset times were estimated to determine day vs. night flight proportions like in other species.

For both aerial feeders and non-stop flyers, we acknowledge that this manual labeling can introduce more observer bias than in other species, so we did not calculate precise departure or landing times. Still, our estimates clearly show multiple partially or fully nocturnal migratory flights, a behavior previously unknown or poorly documented in these species.^{31,65} Moreover, even with a latitudinal error margin of ~ 1000 km, sunset variation is minimal (0–2 min in September; 5–9 min in October) across West Africa (5–15° N), making our nocturnal/diurnal flight proportions reasonably robust at the species level.

Ecological barrier crossing flights

Ecological barriers are natural geographic features or environmental conditions that significantly hinder the migration of species. For migratory animals, these barriers can restrict access to suitable habitats, increase travel risks, and influence migration routes and timing.^{66,67} When crossing large marine expanses, migratory landbirds have no opportunity to land. If the crossing takes longer than a single night, part of the flight must necessarily occur during the day. To enable consistent comparisons across species and populations, some of which may not face such ecological barriers, we also estimated nocturnal and diurnal flight times excluding flights that involved barrier crossings. We focused on two key ecological barriers: marine expanses over 500 km and the Sahara Desert.

The 500 km marine threshold was informed by tracking data from five northern wheatears *Oenanthe oenanthe* undertaking their first autumnal flight from Ouessant island to northwest Iberia. These birds took an average of 11.1 hours to complete the ~ 500 km journey, departing at civil dusk and arriving a few hours after civil dawn.²⁷ This suggests that 500 km is close to the upper limit of what small passerines can fly overnight without needing to continue into the day. Therefore, we used 500 km as a conservative threshold beyond which flights starting in the evening are likely to involve daytime movement. We chose northern wheatear tracking data specifically because they provided empirically grounded estimates of overnight flight distance for a small passerine undertaking a well-defined sea crossing, rather than relying on theoretical flight range estimates. This threshold is conservative as species capable of longer overnight flights would require a greater crossing distance before daytime prolongation becomes necessary.

To classify a flight as crossing an ecological barrier, we considered whether at least half of its duration occurred within the defined polygon representing that barrier. This criterion ensures that the majority of the flight is meaningfully associated with the barrier environment, while also accounting for uncertainties in trajectory estimation and barrier boundaries. We used a global polygon of all marine expanses and a polygon representing the boundaries of the Sahara Desert as defined in.⁶⁸

QUANTIFICATION AND STATISTICAL ANALYSIS

Phylogenetic modelling

We generated a maximum clade credibility phylogenetic tree including all 56 bird species examined in this study using the *clootl* R package,⁵⁶ which accesses the Cornell Lab of Ornithology Open Tree of Life Avian Phylogeny (version Aves v1.4).⁶⁹ It represents one of the most up-to-date and comprehensive avian phylogenetic frameworks currently available, synthesizing phylogenetic information from all recently published studies (see⁶⁹).

We used four indices to evaluate the phylogenetic signal in the proportion of nocturnal flight time, whether including or excluding ecological barrier crossing flights: Pagel's λ , Blomberg's K, Abouheif's Cmean and Moran's I index.⁷⁰ Overall, values of λ , K, I and Cmean larger than zero denote deep relationships between species' traits and phylogeny.⁷⁰ Under a Brownian Motion (BM) model of trait evolution, Pagel's λ and Blomberg's K are expected to be equal to 1. When values of λ and K are close to 0, there is no phylogenetic signal in the studied trait, that is, there is phylogenetic independence, and the trait has evolved independently of phylogeny and close relatives are not more similar than distant relatives. Here, we used Blomberg's K*, a modification of the original Blomberg's K (see Blomberg and Garland⁷¹) that is more robust across different tree shapes and trait distributions. The I and Cmean are autocorrelation indices and measure how similar trait values are between neighboring species on the phylogeny; they are not based on any evolutionary model and large deviations from 0 are representative of strong phylogenetic signal.⁷⁰ We evaluated all four metrics because they are complementary and capture different aspects of phylogenetic signals. We used the *phylosignal* R package to run these analyses.⁵⁸

To assess whether the phylogenetic structuring of nocturnal migration could be attributed to shared ecological conditions among related species rather than true evolutionary conservatism, we ran phylogenetic generalized least squares models (PGLS) using the *nlme* R package.⁵⁷ We used the species-level mean proportion of nocturnal flights (including barrier-crossing flights) as the response variable, and breeding latitude and total migration distance as ecological covariates (see Figure S4). Breeding latitude was computed as the mean absolute latitude of capture locations per species from our tracking dataset. Total migration distance was estimated as the cumulative great-circle distance between consecutive stopover locations along the most likely trajectory of each individual, averaged per species, using the *geosphere* R package.⁵⁹