

Short communication

Hyalomma rufipes of Asian origin transported to Europe by a migrant birdAttila D. Sándor^{a,b,c,*}, Áron Péter^b, Joanna B. Wong^d, Reto Burri^d, Sándor Hornok^{a,b}^a HUN-REN-UVMB Climate Change: New Blood-Sucking Parasites and Vector-Borne Pathogens Research Group, Hungary^b Department of Parasitology and Zoology, University of Veterinary Medicine, Budapest, Hungary^c STAR-UBB Institute, Babes-Bolyai University, Cluj-Napoca, Romania^d Swiss Ornithological Institute, CH-6204 Sempach, Switzerland

ARTICLE INFO

Keywords:

Bird migration
Ixodidae
Parasite transfer
Ticks
Vector

ABSTRACT

Hyalomma rufipes is one of the most epidemiologically important ticks in Africa and the Middle East. It is regularly transported by migratory birds and there are chances that these ticks may become sources for the emergence of resident populations in the temperate region due to recent changes in climatic conditions. In May 2023, a *Hyalomma* sp. nymph was collected in SE Romania from a long-distance migrant host, the Pied Wheatear (*Oenanthe pleschanka*) with known migratory route. The tick was identified morphologically and genetically as *H. rufipes* and based on the timing of the migratory track it attached to its avian host in Saudi Arabia, the Middle East. This is the first ever *H. rufipes* recorded in Europe with known Middle Eastern/Asian origin.

1. Introduction

Ticks are currently the most important vectors of zoonotic diseases in the Northern Hemisphere, with several tick-borne diseases showing emerging patterns (Madison-Antenucci et al., 2020). Bacterial diseases like Lyme borreliosis, human granulocytic anaplasmosis, or relapsing fevers are on the rise both in Europe, as well as in North America, while the impact of viral tick-borne encephalitis or Crimean-Congo Haemorrhagic fever (CCHF) seems to be vastly underestimated both in Africa and Asia (Heyman et al., 2010; Mackenzie and Williams, 2009; Temur et al., 2021). All these emerging health risks have a common background: the changes in geographic occurrences of their main vectors, the ticks. Several epidemiologically important tick species were documented to expand ranges, with northward or elevational shifts, or new colonisations (Cunze et al., 2022; Medlock et al., 2013). Among these tick species, one bears particular importance, *Hyalomma rufipes* Koch, 1844, locally also known as “the hairy *Hyalomma*” or “the coarse bont-legged *Hyalomma*” (Latif, 2013; Walker, 2003).

Hyalomma rufipes is a two-host tick species, which was considered a subspecies of *Hyalomma marginatum* for a long period (Horak et al., 2002), but it is currently classified as a valid species (Apanaskevich and Horak, 2008). It is primarily an African species, occurring in more arid habitats, with a distribution split in two by the tropical areas. It does occur outside Africa, with resident populations in some countries in the

Middle East (Yemen, Oman and Saudi Arabia), but its sporadic occurrence was also reported from most Mediterranean countries, and Central Europe east to Caucasus, Russia, Iraq and China (Vatansever, 2017). *Hyalomma rufipes* is the most common vector of Crimean-Congo Haemorrhagic Fever (CCHF) virus in Africa, while blamed for the transmission of several *Rickettsia* species, *Theileria annulata*, but also *Babesia occultans* (Bonnet et al., 2023), thus making it a very important tick species both from a human epidemiologic, as well as a livestock health perspective. Adult ticks rely on large herbivores (mainly cattle, horses, sheep and camels, but also wild ungulates and ostriches) for feeding, while both larvae and nymphs use small mammals, reptiles and birds as hosts (Vatansever, 2017). As such, the immatures of the species regularly occur on Palearctic-African long-distance migrant birds, which may transfer these ticks from Africa, all along their northward spring migration (Hoogstraal and Kaiser, 1958). Although larvae and mostly nymphs were regularly recorded on such migrants in several countries in Europe (Keve et al., 2022), the occurrence of adult *H. rufipes* in Europe is rare and mostly confined to single individuals found mainly on horses (Chitimia-Dobler et al., 2016). An establishment of permanent populations in Europe was considered unlikely due to climatic conditions, however, there are signs of changes in this situation with a small, reproductive population recorded in Hungary (Keve et al., 2023). The origin of these ticks was hypothesised to be of sub-Saharan Africa, with all molecularly tested, European *H. rufipes* records showing high

* Corresponding author at: HUN-REN Climate Change: New Blood-Sucking Parasites and Vector-Borne Pathogens Research Group, István u. 2, Budapest, Hungary.
E-mail addresses: Sandor.Atila@univet.hu (A.D. Sándor), peter.aron@usamvcluj.ro (Á. Péter), joanna.wong@vogelwarte.ch (J.B. Wong), reto.burri@vogelwarte.ch (R. Burri), Hornok.Sandor@univet.hu (S. Hornok).

<https://doi.org/10.1016/j.ttbdis.2025.102518>

Received 5 May 2025; Received in revised form 23 June 2025; Accepted 5 July 2025

Available online 22 July 2025

1877-959X/© 2025 The Authors. Published by Elsevier GmbH. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

similarity with African sequences (Chitimia-Dobler et al., 2016; Hansford et al., 2019; Hornok et al., 2022; Keve et al., 2023; Lesiczka et al., 2022; Uiterwijk et al., 2021). The aim of the present study was to present a record of a *H. rufipes* with Asian origin collected in Romania from a bird with known migratory route.

2. Materials and methods

The tick was removed from an adult male Pied Wheatear (*Oenanthe pleschanka*) (Fig. 1.) on 05 May 2023 near Mihail Kogalniceanu, SE Romania (44.4182 N; 28.5072E). It was fully engorged and self-detached while the host was processed. For the estimated developmental period of *H. rufipes* subadults, previously published data based on laboratory rearing of ticks were used (Knight et al., 1978; Magano et al., 2000).

After removal, the tick was preserved in 97 % ethanol. The capture of the host bird was a targeted action, as this individual was marked with color rings before and was bearing a miniature data logger. It was initially captured and deployed with the datalogger in the same place in May 2022, to follow its post-breeding migration and wintering. The geolocator used (a multi-sensor 1.2 g GDL3-PAM geolocator, Swiss Ornithological Institute) was mounted using a leg-loop back-harness and collected data on ambient light intensity (Lisovski et al., 2020). Using a preset threshold method (threshold = 1; log-scale), the sunrise and sunset times of each day of the journey was determined from the recorded light data using the R package TwGeos (R Core Team, 2023; Wotherspoon et al., 2016). The tag was calibrated at the breeding site to derive the median reference solar zenith angle, and subsequent location estimation, construction and refinement of the most probable median route (from 2000 final Markov chain Monte Carlo iterations) was modelled using R package SGAT (Wotherspoon et al., 2013).

The collected tick was identified using standard morphological characters (Estrada-Peña et al., 2018). The DNA was extracted from one leg using the QIAamp DNA Mini Kit (QIAGEN, Hilden, Germany), including an overnight digestion in tissue lysis buffer and Proteinase K at 56 °C. PCR amplification of an approx. 710-bp-long fragment of the cytochrome c oxidase subunit I (*cox1*) barcoding gene was attempted with the primer pairs LCO1490 (5'-GGT CAA ATC ATA AAG ATA TTG G-3') and HCO2198 (5'-TAA ACT TCA GGG TGA CCA AAA AAT CA-3') (Folmer et al., 1994). In addition, a second genetic marker, an approx. 460-bp-long fragment of the 16S rDNA gene of Ixodidae was also amplified with the primers 16S+1 (5'-CTG CTC AAT GAT TTT TTA AAT TGC TGT GG-3') and 16S-1 (5'-CCG GTC TGA ACT CAG ATC AAG T-3') (Black and Piesman, 1994). PCR reaction components included 5 µl of

extracted DNA, added to 20 µl of reaction mixture containing 1 U Hot-Star Taq Plus DNA Polymerase (5U/µl) (QIAGEN, Hilden, Germany), 0.5 µl dNTP Mix (10 mM), 0.5 µl of each primer (50 µM), 2.5 µl of 10x Coral Load PCR buffer (15 mM MgCl₂ included) and 15.8 µl ddH₂O. For amplification, an initial denaturation step at 95 °C for 5 min was followed by 40 cycles of denaturation at 94 °C for 40 s, annealing for 1 min at 48 °C or 51 °C in case of the *cox1* and 16S rRNA genes, respectively, and extension at 72 °C for 1 min. Final extension was performed at 72 °C for 10 min. PCR products were visualized in 1.5 % agarose gel.

Purification and sequencing of the PCR products were done by Eurofins Biomi Ltd. (Gödöllő, Hungary). Quality control and trimming of sequences were performed with the BioEdit program. Obtained sequences were compared to GenBank data by the nucleotide BLASTN program (<https://blast.ncbi.nlm.nih.gov>).

3. Results

The collected tick was identified as a fully engorged *Hyalomma* sp. nymph (based on the presence of eyes on the scutum, form and length of hypostome and palps, as well coxa I, see Fig. 2). The *cox I* gene sequence (accession no PV810458) showed a 100 % identity with sequences of *H. rufipes* from Nigeria (GenBank accession no MN601293), Kenya (OQ540949), and Cameroon (MK648422). The 16S rRNA sequence (accession no PV765685) showed 100 % identity to sequences of *H. rufipes* collected in Namibia (KU130461), South Africa (KU130465), and Senegal (KU130457). Considering the 23–28 days as the developmental period of *H. rufipes* subadults, we localised the geographical area of probable presence of the host 23–28 days prior to its recapture. This time period (ca. on dates between 7–11 April 2023) partly overlaps with a minor migratory staging east of Riyadh, in the region of Haradh, SE Saudi Arabia (a region of ca. 50–70 km radius circle with central coordinates at 23.46 N, 48.78 E). Here the bird interrupted its northward migration for eight days in the mentioned period (see Fig. 3.), this being the likely geographical region where *H. rufipes* attached as larvae, which developed into the nymph removed from the adult Pied Wheatear in Romania.

4. Discussion

Long-distance Palearctic-African migrants are known to regularly



Fig. 1. Adult male Pied Wheatear (*Oenanthe pleschanka*) at the moment of initial capture on 27 May 2022.



Fig. 2. Fully engorged nymph of *Hyalomma rufipes*, collected from a Pied Wheatear (*Oenanthe pleschanka*) on 08 May 2023 near Mihail Kogalniceanu, SE Romania.




Fig. 3. Reconstructed migratory track of the adult Pied Wheatear (*Oenanthe pleschanka*), ring no COR R032468, heatmap generated using coordinates based on sunrise/sunset records collected by a geolocator, showing ringing/recapture site (point *a*) as well probable region of tick acquisition (ca. 200 km radius circle *b*, enlarged in the right upper corner showing landscape composition using satellite image).

transport ticks of tropical or subtropical origin to temperate regions of Europe or Asia (Hornok et al., 2022; Keve et al., 2022). Among these migrants, open landscape specialists, like *Oenanthe* spp. wheatears are considered important hosts for *Hyalomma* spp. ticks (Hoogstraal et al., 1961; Hoogstraal and Kaiser, 1958; Kaiser et al., 1974; Keve et al., 2022), with regular records of these species hosting *H. rufipes* in North Africa (Hoogstraal and Kaiser, 1958), Europe (Mancuso et al., 2022; Saikku et al., 1971) or the Middle East (Kaiser et al., 1974). The Pied

Wheatear is a typical representative of this group, with a primarily Asian distribution, and having the most western breeding population along the shores of the Black Sea (Schweizer et al., 2019). These birds winter in East Africa (especially in the Horn of Africa, see Cramp and Perrins, 1993), and spring migrants were already recorded to host nymphs of *H. rufipes* in Cyprus (Kaiser et al., 1974) and in coastal Saudi Arabia (Fain et al., 1995). Based on the migratory route of this tracked individual Pied Wheatear, the most likely point of origin of the nymphal tick

lays in the SE part of Saudi Arabia, east to Riyadh, in the region of Haradh. Unfortunately, with the considerable uncertainty of geolocator provided data (Lisovski and Hahn, 2012), we are not able to pinpoint the exact location the bird used. However, this whole region shows fairly similar habitat pattern of extensive areas of sandy-gravel desert, with rocks and salt lakes (sabkha-s) to the south with low vegetation cover and ideal habitats for larks and wheatears (Boland et al., 2020). The region is a hotspot of Pied Wheatear records in Saudi Arabia. The largest integrated dairy farm in the world is also located here, with extensive arable and grazing fields, providing suitable conditions for ticks, like *H. rufipes*, which is a common parasite of livestock in the region (Abdally et al., 2020; Alghamdi, 2024).

To the best of our knowledge, this is the first geolocated record of any tick of Middle Eastern/Asian origin in Europe and it is also the first observation of *H. rufipes* in Romania. Immatures of *H. rufipes* are commonly found on many bird species (52 species listed by Keve et al., 2022), and it is known that such tick individuals may harbour pathogens, too. Molecular analyses of bird-derived ticks registered DNA of several rickettsial species in *H. rufipes* nymphs, with *Rickettsia aeschlimannii* being the most common (Battisti et al., 2020; Mancuso et al., 2022; Olivieri et al., 2021). However, this tick is the most important vector for CCHF virus in Africa and it is involved in the circulation of two apicomplexans with huge impact on livestock health, *Babesia occultans* (Decaro et al., 2013) and *Theileria annulata* (Bonnet et al., 2023). While neither of these pathogens were recorded in Romania up to now, serological screening of livestock showed 27–37 % seroprevalence for CCHF virus in SE Romania (Bratuleanu et al., 2022; Ceianu et al., 2012), suggesting at least occasional presence of this pathogen. As current changes in land use and climatic conditions may favour the establishment of new vector species in non-endemic areas, the screening of migratory birds may be used as an early warning system to highlight such colonisations and to trigger timely response in case of any new pathogen introduction by these vectors.

Author statement

Hereby the authors declare that they have not used any type of generative artificial intelligence for the writing of this manuscript, nor for the creation of images, graphics, tables, or their corresponding captions.

Ethics approval and consent to participate

Birds were captured and tagged during routine bird ringing operations required for the study of population dynamics of migratory birds according the national legislation (Law 49/2011, Annex 8, art. C), number of ringing permit COR828265/2001 (issued to ADS).

Consent for publication

Not applicable.

CRediT authorship contribution statement

Attila D. Sándor: Writing – review & editing, Writing – original draft, Investigation, Data curation, Conceptualization. **Aron Péter:** Writing – review & editing, Visualization, Resources, Investigation, Data curation. **Joanna B. Wong:** Writing – review & editing, Resources, Investigation, Formal analysis, Data curation. **Reto Burri:** Writing – review & editing, Resources, Project administration, Methodology, Funding acquisition, Formal analysis, Data curation. **Sándor Hornok:** Writing – review & editing, Resources, Investigation, Formal analysis, Data curation, Conceptualization.

Declaration of competing interest

The authors declare that they have no competing interests.

Acknowledgements

The authors would like to express their gratitude to the volunteers who took part in fieldwork and Pied Wheatear tagging, namely Niloofar Alaei Kakhki, Gilles Hauser, Zsolt Kovács, Dave Lutgen and Marco Pilati. We also thank the help provided by C. Domşa and B. Sándor in the preparation of figures, while N. Takács contributed with PCRs and A. Corduneanu helped with GenBANK submission. The work of ADS and SH was funded by the Office for Supported Research Groups, Hungarian Research Network (HUN-REN), Hungary (Project No.1500107).

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.ttbdis.2025.102518.

Data availability

All data related to this study is presented in the manuscript.

References

- Abdally, M.H., Al-Marri, T.M., Abdally, H.M., Al-Jabr, O.A., 2020. Incidence and prevalence of hard ticks in ruminants of Al-Ahsa oasis region, Kingdom of Saudi Arabia. *World's Vet. J.* 10, 276–285. <https://doi.org/10.36380/sci.2020.vwj36>.
- Alghamdi, S.Q., 2024. Prevalence of tick infestation and molecular characterization of tick species of camels (*Camelus dromedarius*) from Al-Baha South Saudi Arabia. *J. Umm Al-Qura Univ. Appl. Sci.* 10, 707–717. <https://doi.org/10.1007/s43994-024-00138-w>.
- Apanaskevich, D.A., Horak, I.G., 2008. The genus *Hyalomma* Koch, 1844: v. re-evaluation of the taxonomic rank of taxa comprising the *H. (Euhyalomma) marginatum* Koch complex of species (Acari: ixodidae) with redescription of all parasitic stages and notes on biology. *Int. J. Acarol.* 34, 13–42. <https://doi.org/10.1080/01647950808683704>.
- Battisti, E., Urach, K., Hodžić, A., Fusani, L., Hufnagl, P., Felsberger, G., Ferroglio, E., Duscher, G.G., 2020. Zoonotic pathogens in ticks from migratory birds, Italy. *Emerg. Infect. Dis.* 26, 2986. <https://doi.org/10.3201/eid2612.181686>.
- Black, W.C., Piesman, J., 1994. Phylogeny of hard- and soft-tick taxa (Acari: ixodida) based on mitochondrial 16S rDNA sequences. *Proc. Natl. Acad. Sci. U.S.A.* 91, 10034–10038. <https://doi.org/10.1073/pnas.91.21.10034>.
- Boland, C., Abdullah, A., Babbington, J., Shobrak, M., Al-Boug, A., 2020. The birds of Saudi Arabia. Saudi Aramco.
- Bonnet, S.I., Bertagnoli, S., Falchi, A., Fígoni, J., Fite, J., Hoch, T., Quillery, E., Moutailler, S., Raffetin, A., René-Martellet, M., 2023. An update of evidence for pathogen transmission by ticks of the genus *Hyalomma*. *Pathogens* 12, 513. <https://doi.org/10.3390/pathogens12040513>.
- Bratuleanu, B., Anita, A., Temmam, S., Dascalu, A., Crivei, L., Cozma, A., Pourquier, P., Savuta, G., Eloit, M., Anita, D., 2022. Seroprevalence of Crimean-Congo hemorrhagic fever among small ruminants from southern Romania. *Vector-Borne Zoo. Dis.* 22, 397–401. <https://doi.org/10.1089/vbz.2021.0091>.
- Ceianu, C.S., Panculescu-Gatej, R.I., Coudrier, D., Bouloy, M., 2012. First serologic evidence for the circulation of Crimean-Congo Hemorrhagic Fever Virus in Romania. *Vector-Borne Zoonotic Dis.* 12, 718–721. <https://doi.org/10.1089/vbz.2011.0768>.
- Chitimia-Dobler, L., Nava, S., Bestehorn, M., Dobler, G., Wölfel, S., 2016. First detection of *Hyalomma rufipes* in Germany. *Ticks Tick-borne Dis.* 7, 1135–1138. <https://doi.org/10.1016/j.ttbdis.2016.08.008>.
- Cramp, S., Perrins, C.M., 1993. *Handbook of the Birds of Europe the Middle East and North Africa. The Birds of the Western Palearctic*. Oxford Univ. Press.
- Cunze, S., Glock, G., Kochmann, J., Klimpel, S., 2022. Ticks on the move—climate change-induced range shifts of three tick species in Europe: current and future habitat suitability for *Ixodes ricinus* in comparison with *Dermacentor reticulatus* and *Dermacentor marginatus*. *Parasitol. Res.* 121, 2241–2252. <https://doi.org/10.1007/s00436-022-07556-x>.
- Decaro, N., Larocca, V., Parisi, A., Losurdo, M., Lia, R.P., Greco, M.F., Miccolis, A., Ventrella, G., Otranto, D., Buonavoglia, C., 2013. Clinical bovine piroplasmiasis caused by *Babesia occultans* in Italy. *J. Clin. Microbiol.* 51, 2432–2434. <https://doi.org/10.1128/JCM.00713-13>.
- Estrada-Peña, A., Mihalca, A.D., Petney, T., 2018. *Ticks of Europe and North Africa: a Guide to Species Identification*. Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-319-63760-0>.
- Fain, A., Roggeman, W., Symens, P., Al Suhaibani, A., 1995. Notes on a small collection of immature *Hyalomma* ticks from migrating birds during their spring migration in Karan Island, Persian Gulf, Saudi Arabia. *Bull. Inst. Roy. Sci. Nat. Belg. Entomol.* 65, 101–103.

- Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R., 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* 3, 294–299.
- Hansford, K.M., Carter, D., Gillingham, E.L., Hernandez-Triana, L.M., Chamberlain, J., Cull, B., McGinley, L., Phipps, L.P., Medlock, J.M., 2019. *Hyalomma rufipes* on an untraveled horse: is this the first evidence of *Hyalomma* nymphs successfully moulting in the United Kingdom? *Ticks Tick-borne Dis.* 10, 704–708. <https://doi.org/10.1016/j.ttbdis.2019.03.003>.
- Heyman, P., Cochez, C., Hoffhuis, A., Van Der Giessen, J., Sprong, H., Porter, S.R., Losson, B., Saegerman, C., Donoso-Mantke, O., Niedrig, M., Papa, A., 2010. A clear and present danger: tick-borne diseases in Europe. *Expert Rev. Anti-Infect. Ther.* 8, 33–50. <https://doi.org/10.1586/eri.09.118>.
- Hoogstraal, H., Kaiser, M.N., 1958. Observations on Egyptian *Hyalomma* ticks (Ixodoidea, Ixodidae). 2. Parasitism of migrating birds by immature *H. rufipes* Koch. *Ann. Entomol. Soc. Am.* 51, 12–16. <https://doi.org/10.1093/aesa/51.1.12>.
- Hoogstraal, H., Kaiser, M.N., Traylor, M.A., Gaber, S., Guindy, E., 1961. Ticks (Ixodoidea) on birds migrating from Africa to Europe and Asia. *Bull. World Health Organ.* 24, 197–212.
- Horak, I.G., Camicas, J.-L., Keirans, J.E., 2002. The Argasidae, Ixodidae and Nuttalliellidae (Acari: ixodida): a world list of valid tick names. *Exp. Appl. Acarol.* 28, 27–54. <https://doi.org/10.1023/A:1025381712339>.
- Hornok, S., Cutajar, B., Takács, N., Galea, N., Attard, D., Coleiro, C., Galea, R., Keve, G., Sándor, A.D., Kontschán, J., 2022. On the way between Africa and Europe: molecular taxonomy of ticks collected from birds in Malta. *Ticks Tick-borne Dis.* 13, 102001. <https://doi.org/10.1016/j.ttbdis.2022.102001>.
- Kaiser, M.N., Hoogstraal, H., Watson, G.E., 1974. Ticks (Ixodoidea) on migrating birds in Cyprus, fall 1967 and spring 1968, and epidemiological considerations. *Bull. Entomol. Res.* 64, 97–110. <https://doi.org/10.1017/S0007485300027024>.
- Keve, G., Csörgő, T., Benke, A., Huber, A., Mórocz, A., Németh, Á., Kalocs, B., Tamás, E.A., Gyurácz, J., Kiss, O., 2023. Ornithological and molecular evidence of a reproducing *Hyalomma rufipes* population under continental climate in Europe. *Front. Vet. Sci.* 10, 1147186. <https://doi.org/10.3389/fvets.2023.1147186>.
- Keve, G., Sándor, A.D., Hornok, S., 2022. Hard ticks (Acari: ixodidae) associated with birds in Europe: review of literature data. *Front. Vet. Sci.* 9, 928756. <https://doi.org/10.3389/fvets.2022.928756>.
- Knight, M.M., Norval, R.A.I., Rechav, Y., 1978. The life cycle of the tick *Hyalomma marginatum rufipes* Koch (Acarina: ixodidae) under laboratory conditions. *J. Parasitol.* 143–146. <https://doi.org/10.2307/3279627>.
- Latif, A.A., 2013. Ixodid ticks of major economic importance and their distribution in South Africa. *Ticks and Tick-Borne Diseases Monograph* 1. AGRI Connect (PTY) LTD.
- Lesiczka, P.M., Daněk, O., Modrý, D., Hrazdilová, K., Votýpka, J., Zurek, L., 2022. A new report of adult *Hyalomma marginatum* and *Hyalomma rufipes* in the Czech Republic. *Ticks Tick-borne Dis.* 13, 101894. <https://doi.org/10.1016/j.ttbdis.2021.101894>.
- Lisovski, S., Bauer, S., Briedis, M., Davidson, S.C., Dhanjal-Adams, K.L., Hallworth, M.T., Karagicheva, J., Meier, C.M., Merkel, B., Ouwehand, J., Pedersen, L., Rakhimberdiev, E., Roberto-Charon, A., Seavy, N.E., Sumner, M.D., Taylor, C.M., Wotherspoon, S.J., Bridge, E.S., 2020. Light-level geolocator analyses: a user's guide. *J. Anim. Ecol.* 89, 221–236. <https://doi.org/10.1111/1365-2656.13036>.
- Lisovski, S., Hahn, S., 2012. GeoLight – processing and analysing light-based geolocator data in R. *Methods Ecol. Evol.* 3, 1055–1059. <https://doi.org/10.1111/j.2041-210X.2012.00248.x>.
- Mackenzie, J.S., Williams, D.T., 2009. The zoonotic flaviviruses of southern, south-eastern and eastern Asia, and Australasia: the potential for emergent viruses. *Zoonoses. Public Health* 56, 338–356. <https://doi.org/10.1111/j.1863-2378.2008.01208.x>.
- Madison-Antenucci, S., Kramer, L.D., Gebhardt, L.L., Kauffman, E., 2020. Emerging tick-borne diseases. *Clin. Microbiol. Rev.* 33. <https://doi.org/10.1128/CMR.00083-18.e00083-18>.
- Magano, S.R., Els, D.A., Chown, S.L., 2000. Feeding patterns of immature stages of *Hyalomma truncatum* and *Hyalomma marginatum rufipes* on different hosts. *Exp. Appl. Acarol.* 24, 301–313. <https://doi.org/10.1023/A:1006457325097>.
- Mancuso, E., Toma, L., Pascucci, I., d'Alessio, S.G., Marini, V., Quaglia, M., Riello, S., Ferri, A., Spina, F., Serra, L., 2022. Direct and indirect role of migratory birds in spreading CCHFV and WNV: a multidisciplinary study on three stop-over islands in Italy. *Pathogens* 11, 1056. <https://doi.org/10.3390/pathogens11091056>.
- Medlock, J.M., Hansford, K.M., Bormane, A., Derdakova, M., Estrada-Peña, A., George, J.-C., Golovljova, I., Jaenson, T.G.T., Jensen, J.-K., Jensen, P.M., Kazimirova, M., Oteo, J.A., Papa, A., Pfister, K., Plantard, O., Randolph, S.E., Rizzoli, A., Santos-Silva, M.M., Sprong, H., Vial, L., Hendrickx, G., Zeller, H., Van Bortel, W., 2013. Driving forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. *Parasites Vect.* 6, 1. <https://doi.org/10.1186/1756-3305-6-1>.
- Olivieri, E., Kariuki, E., Floriano, A.M., Castelli, M., Tafesse, Y.M., Magoga, G., Kumsa, B., Montagna, M., Sasser, D., 2021. Multi-country investigation of the diversity and associated microorganisms isolated from tick species from domestic animals, wildlife and vegetation in selected african countries. *Exp. Appl. Acarol.* 83, 427–448. <https://doi.org/10.1007/s10493-021-00598-3>.
- R Core Team, 2023. R: A language and Environment For Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Saikkku, P., Ulmanen, I., Brummer-Korvenkontio, M., 1971. Ticks (Ixodidae) on migratory birds in Finland. *Acta Entomol. Fen.* 28, 46–51.
- Schweizer, M., Warmuth, V., Alaei Kakhki, N., Aliabadian, M., Förschler, M., Shirihi, H., Suh, A., Burri, R., 2019. Parallel plumage colour evolution and introgressive hybridization in wheatears. *J. Evol. Biol.* 32, 100–110. <https://doi.org/10.1111/jeb.13401>.
- Temur, A.I., Kuhn, J.H., Pecor, D.B., Apanaskevich, D.A., Keshtkar-Jahromi, M., 2021. Epidemiology of Crimean-Congo hemorrhagic fever (CCHF) in Africa—Underestimated for decades. *Am. J. Trop. Med. Hyg.* 104, 1978–1990. <https://doi.org/10.4269/ajtmh.20-1413>.
- Uiterwijk, M., Ibáñez-Justicia, A., Van De Vossenberg, B., Jacobs, F., Overgaaauw, P., Nijse, R., Dabekausen, C., Stroo, A., Sprong, H., 2021. Imported *Hyalomma ticks* in The Netherlands 2018–2020. *Parasit. Vect.* 14, 244. <https://doi.org/10.1186/s13071-021-04738-x>.
- Vatansever, Z., 2017. *Hyalomma rufipes* Koch, 1844 (Figs. 142–144). In: Estrada-Peña, A., Mihalca, A.D., Petney, T.N. (Eds.), *Ticks of Europe and North Africa*. Springer International Publishing, Cham, pp. 355–359. https://doi.org/10.1007/978-3-319-63760-0_67.
- Walker, A.R., 2003. Ticks of domestic animals in Africa: a guide to identification of species. Bioscience reports Edinburgh.
- Wotherspoon, S., Sumner, M., Lisovski, S., 2016. TwGeos: basic data processing for light-level geolocation archival tags.
- Wotherspoon, S.J., Sumner, M.D., Lisovski, S., 2013. R package SGAT: solar/satellite geolocation for animal tracking. GitHub Repository-<http://github.com/swotherspoon/sgat>.